



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ


**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.


In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *Andrew Gersey*

Dated 15 October 2003



PCT/EP 03 / 1 2 1 9 1

25 JUL 03 E825211-2 002029

P01/7700 0.00-0317380.4

Patents Form 1/77

Patents Act 1977

(Rule 16)

24 JUL 2003

The
Patent

Office

24 JUL 2003

LONDON

1/77

Request for grant of a patent

(See the notes on the back of this form. You can also get
an explanatory leaflet from the Patent Office to help
you fill in this form)

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

CJ/KMM/VB60410P

2. Patent application number

(The Patent Office will fill in his part)

0317380.4

3. Full name, address and postcode of the or of
each applicant (underline all surnames)

GlaxoSmithKline Biologicals s.a.
Rue de l'Institut 89, B-1330 Rixensart, , Belgium

Patents ADP number (*if you know it*)

Belgian

If the applicant is a corporate body, give the
country/state of its incorporation

08101271001

4. Title of the invention

Drying Method

5. Name of your agent (*if you have one*)

Corporate Intellectual Property

"Address for service" in the United Kingdom
to which all correspondence should be sent

(including the postcode)

08072555006

Patents ADP number (*if you know it*)

GlaxoSmithKline
Corporate Intellectual Property (CN9 25.1)
980 Great West Road
BRENTFORD
Middlesex TW8 9GS

6. If you are declaring priority from one or more
earlier patent applications, give the country
and the date of filing of the or each of
these earlier applications and (*if you know it*) the
or each application number

Country	Priority application number (<i>if you know it</i>)	Date of filing (<i>day / month / year</i>)
---------	--	---

7. If this application is divided or otherwise
derived from an earlier UK application,
give the number and the filing date of
the earlier application

Number of earlier application	Date of filing (<i>day / month / year</i>)
-------------------------------	---

8. Is a statement of inventorship and of right

Drying Process

The present invention relates to the preservation of biological and other labile samples, to such preserved samples and to a novel process for preserving such samples. The novel process comprises adding a sample including an active agent and a stabilising agent to a container, subjecting the sample to such temperature and pressure conditions to cause solvent loss by evaporation without freezing the sample or bubbling to form a foam. Subsequently, during a secondary drying phase, pressure and temperature conditions are maintained or adjusted so that solvent is removed and the preservation sample dries to form a highly viscous liquid. Further provided by the present invention are compositions preserved by the process of the present invention and in particular preserved vaccine compositions.

There is a need to extend the stability and thus the shelf life of labile samples, particularly biological samples. Traditionally, this has been accomplished using the process of freeze drying in which a solution of the substance is made and the sample is frozen. During the primary drying phase, most of the water is removed by sublimation from ice under reduced pressure conditions and a porous 'cake' is formed. This is usually followed by a secondary drying phase when the pressure and temperature are changed and water is evaporated from the solid 'cake'. The resulting lyophilised sample has improved stability compared to a liquid formulation. However, the freeze drying process is lengthy, expensive and can be the rate limiting step in a production process.

Freeze drying can also lead to the loss of activity or antigenicity of some active agents. For certain biological materials such as live virus, there can be significant loss of activity during the freeze drying process (Pikal (1994) ACS Symposium 567: 120-133). Many freeze dried substances are still unstable at ambient temperature (Carpenter et al (1994) ACS Symposium 567; 134-147).

Damage caused by the process of freezing may be circumvented to some degree by the use of stabilising agents such as polyols. Further improvements on the process of

lyophilisation have also been made by avoiding freezing the sample during the process and removing water by boiling (WO96/40077; US6306345). This method involves preparing a mixture of a glass-matrix forming material in a suitable solvent together with the sample to be preserved, evaporating bulk solvent from the mixture to obtain a syrup, exposing the syrup to a pressure and temperature sufficient to cause boiling of the syrup and removing residual solvent. Methods similar to this may be referred to as foam drying techniques. Such techniques will expose the sample to be preserved to stresses due to the formation and bursting of bubbles during the 'boiling' stage. Especially where labile substances are to be preserved, this may result in a loss of activity.

A similar method was described in US5,766,520, in which the process involves partially removing the water to form a viscous fluid and further subjecting the syrup to vacuum to cause it to 'boil' and further drying at temperatures substantially lower than 100 °C. This method still suffers from some of the problems of conventional freeze-drying. When the process is carried out in a large freeze-dryer, samples will dry at different rates depending on their position on the shelf and this leads to different samples losing different amount of activity during the drying process. This leads to a lack of consistency within a batch.

Trehalose is a polyol that is favoured for its stabilising properties. Trehalose is a naturally occurring, inert, non-reducing and non-toxic, glass-forming disaccharide that was initially found to be associated with the prevention of desiccation damage in some plants and animals. Trehalose is useful in preventing denaturation of a wide variety of substances including proteins, viruses and foodstuffs during desiccation and subsequent storage partly because it has a relatively high glass transition temperature (ca 120 °C in the anhydrous state) (US4891319; US5149653; US5026566). Trehalose also stabilises enzymes (Argall and Smith (1993) *Biochem. Mol. Biol. Int.* 30; 491). Trehalose and a wide variety of stabilising polyols have also been found to be useful in improving the preservation of freeze-dried samples, especially in cases where the sample is prone to loss of activity during the freeze-drying process. Other sugars useful in lyophilisation techniques include sucrose and lactose.

The present invention relates to an improved method of preserving an active agent, particularly if the active agent is labile and prone to loss of activity during a more conventional drying process. The process comprises the steps of preparing a
5 preservation sample by dissolving/suspending an active agent in a solution of a stabilising agent; subjecting the preservation sample to such temperature and pressure conditions that the preservation sample loses solvent by evaporation, without the sample freezing or bubbling to form a foam; and removing solvent until the preservation sample dries to form a highly viscous liquid.

10 The process is very gentle and does not expose the active agent to freezing or boiling and is therefore advantageous over conventional freeze drying and foam drying techniques which would subject the sample to one or both of these stresses. Where the active agent to be preserved is labile, the use of the method of the invention leads to
15 increased retention of activity and/or antigenicity. This can be measured by reconstituting the dried active agent in solvent, preferably water or an aqueous solution, and measuring the activity or antigenicity by a standard assay (for example by ELISA) and comparing the results with that obtained with either an undried sample or with samples dried by freeze drying or foam drying techniques, and then
20 reconstituted.

It is particularly advantageous to dry IPV (inactivated polio virus, the active ingredient in injectable polio vaccine) using the process of the invention. IPV is present in known vaccines as a liquid formulation (WO99/48525). Problems have
25 arisen on attempting to use a solid formulation of IPV in a vaccine since standard freeze drying procedures lead to a loss of IPV antigenicity. The process of the invention leads to much higher integrity of the polio virus antigens, due partially to the reduced time required by the process of the invention. It is advantageous to formulate IPV with together with Hib, preferably as a solid formulation, because the
30 presence of IPV can reduce the amount of interference associated with combining Hib with other antigens.

The process of the invention is advantageous over normal freeze drying since the running cycle is shorter and requires less refrigeration making it more energy efficient. Since the drying process is often the rate limiting step of a process, the use of the method of the invention leads to higher levels of production at reduced expense.

Description of figures

Figure 1 – Photograph of the high viscosity liquid in inverted vials.

Detailed description

The method of the invention is used for preserving an active agent and comprises the steps of:

- preparing a preservation sample by suspending or dissolving an active agent in a solution of a stabilising agent;
- subjecting the preservation sample to such temperature and pressure conditions that the preservation sample loses solvent by evaporation, without freezing or bubbling to form a foam, to form a viscous liquid;
- removing solvent until the sample dries to form a highly viscous liquid.

A method of preserving an active agent produces a form of the active agent that is able to withstand extended storage during which the activity and/or antigenicity of the active agent is maintained. Preferably the active agent retains at least 40, 50, 60, 70, preferably 80, 90, 95% of its original activity/antigenicity over a period of at least 3, 6, 9, 12, 24 months storage at 4 °C. Antigenicity can be measured by standard assays for instance by ELISA.

The method is useful for extending the shelf life of labile products which rapidly lose activity when stored in solution or when exposed to freezing or bubbling to form a foam.

A labile product is prone to loss of activity and/ or to loss of antigens following storage in solution and/or freezing and/or subjecting to stresses such as those involved in bubbling during foam formation.

- 5 It is particularly applicable for use where a lower concentration of the glass forming polyol is advantageous and a shorter drying process is preferred.

10 A highly viscous liquid is defined as a material with a solvent content less than or equal to 25, 20, preferably 15, 10, 8, 5, 4, 3, 2 or 1%. The highly viscous liquid has a sufficiently low solvent content such that the active agent is preserved in a stable state for at least 3, 6, 9, 12 or 24 months at 4 °C, allowing the active agent to retain at least 40, 50, 60, preferably 70, 80, 90, 95% of its activity and/or antigenicity over this period. Preferably, the highly viscous liquid has a solid appearance but is a glass and is able to flow very slowly over a period of hours, more preferably days.

15

Preparation of the preservation sample

- 20 Any stabilising agent is suitable for use in the first step of this invention. Suitable materials include, but are not limited to, all polyols, including carbohydrate and non-carbohydrate polyols. Preferably the stabilising polyol enables the active agent to be stored without substantial loss of activity by denaturation, aggregation or other means. Particularly suitable materials include sugars, sugar alcohols and carbohydrate derivatives. Preferably, the glass forming polyol is a carbohydrate or derivatives thereof, including glucose, maltulose, iso-maltulose, lactulose, sucrose, maltose, 25 lactose, iso-maltose, maltitol, lactitol, palatinit, trehalose, raffinose, stachyose, melezitose or dextran, most preferably trehalose, sucrose, sorbitol, raffinose, mannitol, lactose, lactitol or palatinit, most preferably sucrose, sorbitol, lactose or trehalose.
- 30 Bacterial polysaccharides are particularly advantageous for use as a stabilising agent in an immunogenic composition since they can act both as a stabilising agent and an immunogen.

Carbohydrates include, but are not limited to, monosaccharides, disaccharides, trisaccharides, oligosaccharides and their corresponding sugar alcohols, polyhydroxyl compounds such as carbohydrate derivatives and chemically modified carbohydrates, hydroxyethyl starch and sugar copolymers. Both natural and synthetic carbohydrates are suitable for use. Synthetic carbohydrates include, but are not limited to, those which have the glycosidic bond replaced by a thiol or carbon bond. Both D and L forms of the carbohydrates may be used. The carbohydrate may be non-reducing or reducing. Where a reducing carbohydrate is used, the addition of inhibitors of the Maillard reaction is preferred.

Reducing carbohydrates suitable for use in the invention are those known in the art and include, but are not limited to, glucose, maltose, lactose, fructose, galactose, mannose, maltulose and lactulose. Non-reducing carbohydrates include, but are not limited to, non-reducing glycosides of polyhydroxyl compounds selected from sugar alcohols and other straight chain polyalcohols. Other useful carbohydrates include raffinose, stachyose, melezitose, dextran, sucrose, cellibiose, mannobiose and sugar alcohols. The sugar alcohol glycosides are preferably monoglycosides, in particular the compounds obtained by reduction of disaccharides such as lactose, maltose, lactulose and maltulose.

Particularly preferred carbohydrates are trehalose, sucrose, sorbitol, maltitol, lactitol, palatinit and glucopyranosyl-1→6-mannitol.

Amino acids can act as stabilising agents and can be used by themselves and preferably in combination with a polyol. Preferred amino acids include glycine, alanine, arginine, lysine and glutamine although any amino acid, or a combination of amino acids, peptide, hydrolysed protein or protein such as serum albumin can act as a stabilising agent.

The concentration of the stabilising agent used in the process of the invention may be between 1% and 50% weight/volume, preferably 1-5%, 5-10%, 15-20%, 20-

25% or 25-50%, most preferably less than or equal to 15% or 10%. The amounts of stabilising agent required is proportional to the amount of salts present. Therefore, although levels of stabilising agent between 2% and 10% are preferred, higher concentrations of 10% to 25% may be required to dry samples with a high salt content.

Container

Different mixtures and various container shapes and sizes can be processed simultaneously. Ideally, the container size used is sufficient to contain the initial mixture and accommodate the volume of the solid formed thereof. Typically, this is determined by the mass of the glass forming material, the surface area of the container and the conditions of the glass formation. The mass of glass forming material must be sufficient to give viscous syrup which translates practically as a minimal mass per unit area of container surface. This ratio varies from mixture to mixture and container used, but is easily determined empirically by one skilled in the art by following the procedures set forth herein. Any such vials can be used, including Wheaton moulded and tube-cut vials.

The process of the invention preferably uses containers with a solvent repellent, preferably a water repellent interior surface. This is achieved through coating the interior surface with a hydrophobic composition, for instance by siliconisation. Siliconisation is achieved by processes that are well known to those skilled in the art. In one method, the container is siliconised by rising the interior of the container with an emulsion of silicone, followed by processing through an oven at high temperature, typically 350 °C. Alternatively, the water repellent interior surface is achieved by the container being made of a water repellent composition.

The water repellent interior surface of the container makes the dried product of the process easier to reconstitute since less of the water collects on the sides of the container.

Although singular forms may be used herein, more than one glass matrix-forming material, more than one additive, and more than one substance may be present. Effective amounts of these components are easily determined by one skilled in the art.

5 Solution

The solvent into which the stabilising agent and active agent are mixed can be aqueous, organic, or a mixture of both. Sufficient aqueous solvent to dissolve the glass matrix-forming material and sufficient organic solvent to dissolve a hydrophobic substance may be used, allowing the formation of glass incorporating hydrophobic substance(s).

The choice of solvent will depend upon the nature of the material chosen for glass matrix formation, as well as the nature of any additive and/or substance to be incorporated. The solvent should be of a nature and of sufficient volume to effect adequate solubilization of the glass matrix-forming material as well as any additive and/or substance. If the substance is a hydrophilic material, the liquid will preferably be aqueous to avoid any potential loss of activity due to deleterious solvent interactions. Preferably, the aqueous solvent includes any suitable aqueous solvent known in the art, including, but not limited to, water and biological buffer solutions. Preferably, the aqueous solvent is present in an amount of 5 to 98% by volume, more preferably 80-98% by volume, most preferably 85-98% by volume.

The volume of solvent can vary and will depend upon the glass matrix-forming material and the substance to be incorporated as well as any additives. The minimum volume required is an amount necessary to solubilise the various components. However, homogeneously dispersed suspensions of the substance(s) can also be used. Suitable amounts of the components in specific embodiments are easily determinable by those skilled in the art in light of the examples provided herein.

Various additives can be introduced into the preservation sample. A preferred additive is an inhibitor of the Maillard reaction. Preferably, if the substance and/or glass

matrix-forming material contains carbonyl and amino, imino or guanidino groups, the compositions further contain at least one physiologically acceptable inhibitor of the Maillard reaction in an amount effective to substantially prevent condensation of amino groups and reactive carbonyl groups in the composition. The inhibitor of the
5 Maillard reaction can be any known in the art. The inhibitor is present in an amount sufficient to prevent, or substantially prevent, condensation of amino groups and reactive carbonyl groups. Typically, the amino groups are present on the substance and the carbonyl groups are present on the glass matrix forming material, or the converse. However, the amino acids and carbonyl groups may be intramolecular
10 within either the substance or the carbohydrate.

Various classes of compounds are known to exhibit an inhibiting effect on the Maillard reaction and hence to be of use in the compositions described herein. These compounds are generally either competitive or non-competitive inhibitors of the
15 Maillard reaction. Competitive inhibitors include, but are not limited to, amino acid residues (both D and L), combinations of amino acid residues and peptides. Particularly preferred are lysine, arginine, histidine and tryptophan. Lysine and arginine are the most effective. There are many known non-competitive inhibitors. These include, but are not limited to, aminoguanidine and derivatives and
20 amphotericin B. EP-A-0 433 679 also describes suitable Maillard inhibitors which include 4-hydroxy-5, 8-dioxoquinoline derivatives.

It is advantageous to incorporate a coloured dye into the preservation sample in order to allow easier visualisation of the dried product of the method of the invention. This
25 is particularly important during reconstitution to ensure that the highly viscous liquid is thoroughly reconstituted prior to use. Preferably, the coloured dye maintains its colour at a neutral pH and is compatible with injection into a patient. Most preferably the coloured dye is phenol red.

30 Loss of solvent by evaporation (evaporative drying – step b)

The process of the invention involves subjecting the preservation sample to such pressure and temperature conditions so that the preservation sample loses solvent by evaporation, without the sample freezing or bubbling to form a foam.

- 5 The temperature within the preservation sample will, at times, be different from that external to the sample due to the endothermic nature of the evaporation process. References to temperature are to the conditions external to the preservation sample, for instance, where a large industrial freeze dryer is used, to the temperature of the shelf. This usually corresponds to the freeze dryer temperature setting.

10

Optionally a preliminary step of degassing the preservation sample is present in the method of the invention. The pressure is reduced to at or below 200mBars, preferably between 200 and 35mBars, for a period of at least 5 minutes before the pressure is reduced further.

15

- A preferred embodiment of the invention achieves evaporative drying by reducing the pressure while controlling the temperature conditions. The pressure is adjusted to at or below 30, 25, 20, preferably 15, 12, most preferably 10, 8, 7, 6, 5, 4, 3, 2 or 1 mbar; while maintaining the temperature setting at a temperature above 0 °C, preferably of 20 between 5 °C to 37 °C, 5 °C to 10 °C, 10 °C to 15 °C; 15 °C to 20 °C; 20 °C to 25 °C; 25 °C to 30 °C; or 30 °C to 37 °C. These conditions are maintained for at least 1, 2, 3, 4, 5, 8, 10, 12, 16 or 24 hours. In a particularly preferred embodiment, the pressure is maintained above 2mbars where the temperature setting is 15 °C in order to prevent freezing of the sample. Where a higher temperature setting is used, slightly 25 lower pressure is possible without freezing the sample and where a lower temperature setting is used, the pressure should be maintained at the higher level to prevent freezing. Preferably the conditions are maintained for a sufficient period of time so that the evaporation rate has slowed so that the temperature of the sample is approximately the same as that external to the sample.

30

Preferably, the preservation sample does not freeze or bubble/boil to form a foam and loses solvent to form a viscous liquid or a highly viscous liquid.

Removing solvent to form a solid

5 A subsequent stage of the method of the invention involves removing solvent until the preservation sample dries to form a highly viscous liquid.

A highly viscous liquid is defined as a material with a solvent content less than or equal to 25, 20, preferably 15, 10, 8, 5, 4, 3, 2 or 1%. The highly viscous liquid has a sufficiently low solvent content such that the active agent is preserved in a stable state for at least 3, 6, 9, 12 or 24 months at 4 °C, allowing the active agent to retain at least
10 40, 50, 60, preferably 70, 80, 90, 95% of its activity and/or antigenicity over this period. Preferably, the highly viscous liquid has a solid appearance but is a glass and is able to flow very slowly over a period of hours, more preferably days.

In one embodiment of the invention, this is achieved by maintaining the pressure and
15 temperature conditions at those applied in the first evaporative drying stage. For instance, the pressure is maintained at or below 30, 25, 20, preferably 15, 12, most preferably 10, 8, 7, 6, 5, 4, 3, 2 or 1 mbar, while maintaining the temperature setting at a temperature above 0 °C, preferably of between 5 °C to 37 °C, 5 °C to 10 °C, 10 °C to 15 °C; 15 °C to 20 °C; 20 °C to 25 °C; 25 °C to
20 30 °C; or 30 °C to 37 °C. These temperature and pressure conditions are maintained for 1, 2, 3, 4, 5, 6, 8, 10, 12, 18 hours or more in order to obtain a highly viscous liquid with a solvent content less than or equal to 25, 20, preferably 15, 10, 8, 5, 4, 3, 2 or 1%.

25 Another embodiment of the invention increases the temperature setting during solvent removal to a higher temperature setting than that maintained earlier in the process.

This allows the solvent to leave the sample at a quicker rate so that the method of the invention can be completed in a shorter time. For instance, the temperature setting is increased to above 0 °C, more preferably above 20 °C, preferably between 5 °C and
30 37 °C, 5 °C and 10 °C, 10 °C and 20 °C; 20 °C and 30 °C; more preferably 30 °C and 40 °C; more preferably 40 °C and 50 °C; most preferably 50 °C and 60 °C while maintaining the pressure at or below 30, 25, 20, preferably 15, 12, most preferably 10,

8, 7, 6, 5, 4, 3, 2 or 1 mbar. These temperature and pressure conditions are maintained for 1, 2, 3, 4, 5, 6, 8, 10, 12, 18 hours or more in order to obtain a solid with less than or equal to 25, 20, preferably 15, 10, 8, 5, 4, 3, 2 or 1%. This embodiment requires the active agent to be heat stable at the temperature used for the method to be carried out successfully.

A preferred embodiment of the invention reduces the pressure setting during solvent removal (step c) to a lower pressure setting than that used earlier in the process (step b). This allows the solvent to leave the sample at a quicker rate so that the method of the invention can be completed in a shorter time. It also enables a higher proportion of the solvent to be lost. For instance, the pressure setting is set to at or below 7, 6, preferably 5, 4, 3, more preferably 2, 1.5, 1, most preferably 0.8, 0.5, 0.2, 0.1 or 0.05 mbar, while maintaining the temperature at or above 0 °C, preferably between 10 °C and 20 °C; 20 °C and 30 °C; 30 °C and 35 °C or above 40 °C. These temperature and pressure conditions are maintained for 1, 2, 3, 4, 5, 6, 8, 10, 12, 18 hours or more in order to obtain a solid with a solvent content less than or equal to 25, 20, preferably 15, 10, 8, 5, 4, 3, 2 or 1%.

Preferably, steps b) and c) should be completed in a time equal to or less than 18 hours, more preferably 16, 14, 12, most preferably 10 hours.

Active agent

The method of the invention is useful for preserving any active agent however it is particularly useful in the case of labile active agents that lose activity and/or antigenicity during other preservation processes.

The active agent to be preserved using a method of the invention may comprise a biological system selected from the group consisting of cells, subcellular compositions, bacteria, outer membrane vesicle preparations and viruses, virus components or virus like particles. It may also comprise molecules, for instance proteins, peptides, amino acids, polynucleic acids, oligonucleotides, polysaccharides,

oligosaccharides, polysaccharide – protein conjugates, oligosaccharide-protein conjugates.

- Examples of active agents that can be preserved using a method of the invention
- 5 include any bioactive substances such as pharmaceutically effective substances, including, but not limited to, antiinflammatory drugs, analgesics, tranquillisers, antianxiety drugs, antispasmodics, antidepressants, antipsychotics, tranquillisers, antianxiety drugs, narcotic antagonists, antiparkinsonism agents, cholinergic agonists, chemotherapeutic drugs, immunosuppressive agents, antiviral agents, antimicrobial
- 10 agents, appetite suppressants, anticholinergics, antimetotics, antihistaminics, antimigraine agents, coronary, cerebral or peripheral vasodilators, hormonal agents, contraceptives, antithrombotic agents, diuretics, antihypertensive agents, cardiovascular drugs, opioids, and the like.
- 15 Suitable agents also include therapeutic and prophylactic agents. These include, but are not limited to, any therapeutically effective biological modifier. Such substances include, but are not limited to, subcellular compositions, cells, bacteria, outer membrane vesicle preparations, viruses and molecules including but not limited to, lipids, organics, proteins and peptides (synthetic and natural), peptide mimetics,
- 20 hormones (peptide, steroid and corticosteroid), D and L amino acid polymers, oligosaccharides, polysaccharides, nucleotides, oligonucleotides and nucleic acids, including DNA and RNA, protein nucleic acid hybrids, small molecules and physiologically active analogues thereof. Further, the modifiers may be derived from natural sources or made by recombinant or synthetic means and include analogues,
- 25 agonists and homologs.

As used herein “protein” refers also to peptides and polypeptides. Such proteins include, but are not limited to, enzymes, biopharmaceuticals, growth hormones, growth factors, insulin, antibodies, both monoclonal and polyclonal and fragments

30 thereof, interferons, interleukins and cytokines.

Therapeutic nucleic acid-based agents prepared by the methods described herein are also encompassed by the invention. As used herein, "nucleic acids" includes any therapeutically effective nucleic acids known in the art including, but not limited to DNA, RNA, and physiologically active analogues thereof. The nucleotides may
5 encode genes or may be any vector known in the art of recombinant DNA including, but not limited to, plasmids, retroviruses and adeno-associated viruses.

The preservation of substances which are prophylactically active and carriers thereof are further encompassed by the invention. Preferable compositions include
10 immunogens such as vaccines. Vaccines may be for oral administration or may be for injection after reconstitution. Suitable vaccines include, but are not limited to, live and attenuated viruses, nucleotide vectors encoding antigens, live and attenuated bacteria, protein, polysaccharide, oligosaccharide and/or lipopolysaccharide antigens, antigens plus adjuvants and antigens and/or haptens coupled to carriers. Particularly preferred
15 are vaccines effective against diphtheria, tetanus, pertussis, botulinum, cholera, Dengue, Hepatitis A, B, C and E, *Haemophilus influenzae* b, *Streptococcus pneumoniae*, *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, Group B streptococci, Group A streptococci, herpes virus, *Helicobacterium pylori*, influenza, Japanese encephalitis, meningococci A, B,
20 C, Y, W, measles, mumps, papilloma virus, pneumococci, polio virus, inactivated polio virus (IPV – preferably comprising types 1, 2 and 3 as is standard in the vaccine art, most preferably the Salk polio vaccine), rubella, rotavirus, respiratory syncytial virus, Shigella, tuberculosis, yellow fever and combinations thereof. The antigenic component of vaccines may also be produced by molecular biology techniques to
25 produce recombinant peptides or fusion proteins containing one or more portions of a protein derived from a pathogen. For instance, fusion proteins containing an antigen and the B subunit of cholera toxin have been shown to induce an immune response to the antigen. Sanches et al (1989) Proc. Natl. Acad. Sci. USA 86:481-485. Vaccines are particularly suitable for incorporation into the single-dosage composition. They
30 are stable indefinitely under ambient conditions and can be redissolved in sterile diluent immediately before inoculation.

In a preferred embodiment, the immunogenic composition would comprise capsular polysaccharides derived from one or more of serogroups A, C, W-135 and Y of *Neisseria meningitidis*. A further preferred embodiment would comprise capsular polysaccharides derived from *Streptococcus pneumoniae*. The pneumococcal capsular polysaccharide antigens are preferably selected from serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F and 33F (most preferably from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F). A further preferred embodiment would contain the PRP capsular polysaccharides of *Haemophilus influenzae* type b. A further preferred embodiment would contain the Type 5, Type 8, 336 or PNAG capsular polysaccharides of *Staphylococcus aureus*. A further preferred embodiment would contain the Type I, Type II, Type III or PIA capsular polysaccharides of *Staphylococcus epidermidis*. A further preferred embodiment would contain the Type Ia, Type Ic, Type II or Type III capsular polysaccharides of Group B streptococcus. A further preferred embodiment would contain the capsular polysaccharides of Group A streptococcus, preferably further comprising at least one M protein and more preferably multiple types of M protein.

In one embodiment of the invention, the bacterial polysaccharides are full length, being purified native polysaccharides. In an alternative embodiment of the invention, the polysaccharides are sized between 2 and 20 times, preferably 2-5 times, 5-10 times, 10-15 times or 15-20 times, so that the polysaccharides are smaller in size for greater manageability. Oligosaccharides are used in a preferred embodiment. Oligosaccharides typically contain between 2 and 20 repeat units.

Polysaccharide and oligosaccharides may be unconjugated or conjugated as described below.

Combinations of two or more of the above active agents may be preserved using the method of preservation of the invention. Part or all of a vaccine may be preserved using the method of preservation of the invention.

A preferred active agent to be preserved using the process of the invention comprises IPV (an inactivated mixture of polio virus strains). IPV, particularly the type 3 component, is sensitive to conventional freeze drying and foam drying techniques as shown by the loss of antigens following freeze drying or foam drying and subsequent reconstitution.

IPV is defined as inactivated polio virus (preferably comprising types 1, 2 and 3 as is standard in the vaccine art, most preferably the Salk polio vaccine). A vaccine dose of IPV contains 20-80, preferably 40 or 80 D-antigen units of type 1 (Mahoney), 4-16, preferably 8 or 16 D-antigen units of type 2 (MEF-1) and 20-64, preferably 32 or 64 D-antigen units of type 3 (Saukett).

Preferably, the drying process utilised retains the IPV antigens so that an ELISA on the reconstituted highly viscous liquid, using antibodies against polio virus type 1, 2 and/or 3, gives results that are over 40%; 50%, preferably 60%, 70%, more preferably 80% or 90% the level achieved using the undried IPV. In vivo experiments in which the dried formulation, after reconstitution is inoculated into an animal, preferably a mouse may also be used to assess the degree of antigen retention.

Preferably, IPV is combined with one or more of Hib (*Haemophilus influenzae* type b) PRP polysaccharide and/or meningococcal A, C, W and/or Y polysaccharides and/or pneumococcal polysaccharides. Most preferably the active agents comprise, IPV and Hib; IPV and MenC; IPV, Hib and MenC; Hib and MenC; IPV and MenA and C; Hib and Men A and C; IPV, Hib, Men A and C; Hib, Men C and Y; or IPV, Hib, Men C and Y.

The above particularised active agents may also comprise one or more pneumococcal capsular polysaccharides as described below.

In the above compositions where polysaccharides are used, oligosaccharides may also be employed (as defined below).

Although these compositions may be adjuvanted (as described below), they are preferably unadjuvanted or preferably do not comprise aluminium salts.

Preferably the polysaccharides or oligosaccharides are conjugated to a peptide or carrier protein comprising T-helper epitopes (as described below).

Additional components

The preferred combinations, dried by the process of the invention may be combined with other antigens in a combination vaccine which is desiccated or is preferably a liquid formulation which can be used to reconstitute the dried components. Preferred antigens to be combined with the active agents in the paragraph above include one or more of diphtheria toxoid, tetanus toxoid, whole cell pertussis (Pw), acellular pertussis (Pa) (as described below), Hepatitis B surface antigen, pneumococcal polysaccharides, pneumococcal proteins or any of the antigens listed below. Bacterial polysaccharides may be conjugated to a carrier protein such as tetanus toxoid, tetanus toxoid fragment C, diphtheria toxoid, CRM197, pneumolysin, Protein D (US6342224) as described below.

Active agents preserved using the process of the invention may be formulated with capsular polysaccharides derived from one or more of *Neisseria meningitidis*, *Haemophilus influenzae* b, *Streptococcus pneumoniae*, Group A Streptococci, Group B Streptococci, *Staphylococcus aureus* or *Staphylococcus epidermidis*. In a preferred embodiment, the immunogenic composition would comprise capsular polysaccharides derived from one or more of serogroups A, C, W-135 and Y of *Neisseria meningitidis*. A further preferred embodiment would comprise capsular polysaccharides derived from *Streptococcus pneumoniae*. The pneumococcal capsular polysaccharide antigens are preferably selected from serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F and 33F (most preferably from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F). A further preferred embodiment would contain the PRP capsular polysaccharides of *Haemophilus influenzae* type b. A further preferred embodiment would contain the

Type 5, Type 8, 336 or PNAG capsular polysaccharides of *Staphylococcus aureus*. A further preferred embodiment would contain the Type I, Type II, Type III or PIA capsular polysaccharides of *Staphylococcus epidermidis*. A further preferred embodiment would contain the Type Ia, Type Ic, Type II or Type III capsular polysaccharides of Group B streptococcus. A further preferred embodiment would contain the capsular polysaccharides of Group A streptococcus, preferably further comprising at least one M protein and more preferably multiple types of M protein.

In one embodiment of the invention, the bacterial polysaccharides are full length, being purified native polysaccharides. In an alternative embodiment of the invention, the polysaccharides are sized between 2 and 20 times, preferably 2-5 times, 5-10 times, 10-15 times or 15-20 times, so that the polysaccharides are smaller in size for greater manageability. Oligosaccharides are used in a preferred embodiment. Oligosaccharides typically contain between 2 and 20 repeat units:

Such capsular polysaccharides may be unconjugated or conjugated to a carrier protein such as tetanus toxoid, tetanus toxoid fragment C, diphtheria toxoid, CRM197, pneumolysin, Protein D (US6342224). Tetanus toxin, diphtheria toxin and pneumolysin are detoxified either by genetic mutation and/or preferably by chemical treatment.

The polysaccharide conjugate may be prepared by any known coupling technique. For example the polysaccharide can be coupled via a thioether linkage. This conjugation method relies on activation of the polysaccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide may thus be coupled directly or via a spacer group to an amino group on the carrier protein. Preferably, the cyanate ester is coupled with hexane diamine and the amino-derivatised polysaccharide is conjugated to the carrier protein using heterologation chemistry involving the formation of the thioether linkage. Such conjugates are described in PCT published application WO93/15760 Uniformed Services University.

The conjugates can also be prepared by direct reductive amination methods as described in US 4365170 (Jennings) and US 4673574 (Anderson). Other methods are described in EP-0-161-188, EP-208375 and EP-0-477508.

- 5 A further method involves the coupling of a cyanogen bromide activated polysaccharide derivatised with adipic acid hydrazide (ADH) to the protein carrier by Carbodiimide condensation (Chu C. et al Infect. Immunity, 1983 245 256).

Preferred pneumococcal proteins antigens are those pneumococcal proteins which are
10 exposed on the outer surface of the pneumococcus (capable of being recognised by a host's immune system during at least part of the life cycle of the pneumococcus), or are proteins which are secreted or released by the pneumococcus. Most preferably, the protein is a toxin, adhesin, 2-component signal transducer, or lipoprotein of *Streptococcus pneumoniae*, or fragments thereof. Particularly preferred proteins
15 include, but are not limited to: pneumolysin (preferably detoxified by chemical treatment or mutation) [Mitchell *et al.* Nucleic Acids Res. 1990 Jul 11; 18(13): 4010 "Comparison of pneumolysin genes and proteins from *Streptococcus pneumoniae* types 1 and 2.", Mitchell *et al.* Biochim Biophys Acta 1989 Jan 23; 1007(1): 67-72 "Expression of the pneumolysin gene in *Escherichia coli*: rapid purification and
20 biological properties.", WO 96/05859 (A. Cyanamid), WO 90/06951 (Paton et al), WO 99/03884 (NAVA)]; PspA and transmembrane deletion variants thereof (US 5804193 - Briles *et al.*); PspC and transmembrane deletion variants thereof (WO 97/09994 - Briles et al); PsaA and transmembrane deletion variants thereof (Berry & Paton, Infect Immun 1996 Dec;64(12):5255-62 "Sequence heterogeneity of PsaA, a
25 37-kilodalton putative adhesin essential for virulence of *Streptococcus pneumoniae*"); pneumococcal choline binding proteins and transmembrane deletion variants thereof; CbpA and transmembrane deletion variants thereof (WO 97/41151; WO 99/51266); Glyceraldehyde-3-phosphate - dehydrogenase (Infect. Immun. 1996 64:3544); HSP70 (WO 96/40928); PcpA (Sanchez-Beato et al. *FEMS Microbiol Lett* 1998,
30 164:207-14); M like protein, (EP 0837130) and adhesin 18627, (EP 0834568). Further preferred pneumococcal protein antigens are those disclosed in WO 98/18931, particularly those selected in WO 98/18930 and PCT/US99/30390.

Preferred Neisserial proteins to be formulated with the immunogenic composition of the invention include TbpA (WO93/06861; EP586266; WO92/03467; US5912336), TbpB (WO93/06861; EP586266), Hsf (WO99/31132), NspA (WO96/29412), Hap
5 (PCT/EP99/02766), PorA, PorB, OMP85 (also known as D15) (WO00/23595), PilQ (PCT/EP99/03603), PldA (PCT/EP99/06718), FrpB (WO96/31618 see SEQ ID NO:38), FrpA or FrpC or a conserved portion in common to both of at least 30, 50, 100, 500, 750 amino acids (WO92/01460), LbpA and/or LbpB (PCT/EP98/05117; Schryvers et al Med. Microbiol. 1999 32: 1117), FhaB (WO98/02547), HasR
10 (PCT/EP99/05989), lipo02 (PCT/EP99/08315), MltA (WO99/57280) and ctrA (PCT/EP00/00135).

The vaccine may also optionally comprise antigens providing protection against one or more of Diphtheria, tetanus and *Bordetella pertussis* infections. The pertussis
15 component may be killed whole cell *B. pertussis* (Pw) or acellular pertussis (Pa) which contains at least one antigen (preferably two or all three) from PT, FHA and 69kDa pertactin. Typically, the antigens providing protection against Diphtheria and tetanus would be Diphtheria toxoid and tetanus toxoid. The toxoids may chemically inactivated toxins or toxins inactivated by the introduction of point mutations.
20 Alternatively the highly viscous liquid of the invention may be provided as a kit with the highly viscous liquid glass in one container and liquid DTPa or DTPw in another container. The highly viscous liquid is reconstituted with the liquid DTPa or DTPw vaccine (preferably extemporaneously) and administered as a single vaccine. The DTPa or DTPw vaccine typically is adjuvanted at least in part with aluminium
25 hydroxide (for instance Infanrix® and Tritanrix® vaccines of GlaxoSmithKline Biologicals s.a.).

The vaccine may also optionally comprise one or more antigens that can protect a host against non-typeable *Haemophilus influenzae*, RSV and/or one or more antigens that
30 can protect a host against influenza virus.

Preferred non-typeable *H. influenzae* protein antigens include Fimbrin protein (US 5766608) and fusions comprising peptides therefrom (eg LB1 Fusion) (US 5843464 - Ohio State Research Foundation), OMP26, P6, protein D, TbpA, TbpB, Hia, Hmw1, Hmw2, Hap, and D15.

5

Preferred influenza virus antigens include whole, live or inactivated virus, split influenza virus, grown in eggs or MDCK cells, or Vero cells or whole flu virosomes (as described by R. Gluck, Vaccine, 1992, 10, 915-920) or purified or recombinant proteins thereof, such as HA, NP, NA, or M proteins, or combinations thereof.

10

Preferred RSV (Respiratory Syncytial Virus) antigens include the F glycoprotein, the G glycoprotein, the HN protein, the M protein or derivatives thereof.

It should be appreciated that antigenic compositions of the invention may comprise one or more capsular polysaccharide from a single species of bacteria. Antigenic compositions may also comprise capsular polysaccharides derived from one or more species of bacteria.

15

Immunogenic compositions and vaccines

20 A further aspect of the invention includes immunogenic compositions or vaccines comprising the highly viscous liquid of the invention and a pharmaceutically acceptable excipient.

Preferably, the immunogenic composition or vaccine contains an amount of an adjuvant sufficient to enhance the immune response to the immunogen. Suitable adjuvants include, but are not limited to, aluminium salts, squalene mixtures (SAF-1), muramyl peptide, saponin derivatives, mycobacterium cell wall preparations, monophosphoryl lipid A, mycolic acid derivatives, non-ionic block copolymer surfactants, Quil A, cholera toxin B subunit, polphosphazene and derivatives, and immunostimulating complexes (ISCOMs) such as those described by Takahashi et al. (1990) Nature 344:873-875. For veterinary use and for production of antibodies in animals, mitogenic components of Freund's adjuvant can be used.

25

30

As with all immunogenic compositions or vaccines, the immunologically effective amounts of the immunogens must be determined empirically. Factors to be considered include the immunogenicity, whether or not the immunogen will be
5 complexed with or covalently attached to an adjuvant or carrier protein or other carrier, route of administrations and the number of immunising dosages to be administered. Such factors are known in the vaccine art and it is well within the skill of immunologists to make such determinations without undue experimentation.

10 The active agent can be present in varying concentrations in the highly viscous liquid or vaccine of the invention. Typically, the minimum concentration of the substance is an amount necessary to achieve its intended use, while the maximum concentration is the maximum amount that will remain in solution or homogeneously suspended within the initial mixture. For instance, the minimum amount of a therapeutic agent is
15 preferably one which will provide a single therapeutically effective dosage. For bioactive substances, the minimum concentration is an amount necessary for bioactivity upon reconstitution and the maximum concentration is at the point at which a homogeneous suspension cannot be maintained. In the case of single-dosed units, the amount is that of a single therapeutic application. Generally, it is expected
20 that each dose will comprise 1-100ug of protein antigen, preferably 5-50ug and most preferably 5-25ug. Preferred doses of bacterial polysaccharides are 10-20ug, 10-5ug, 5-2.5ug or 2.5-1ug. The preferred amount of the substance varies from substance to substance but is easily determinable by one of skill in the art.

25 Highly viscous liquid comprising an active agent

Another aspect of the invention is a highly viscous liquid comprising an active agent which is preferably obtainable or obtained using a method of the invention. The active agent preferably retains its activity and/or antigenicity following drying using the
30 method of the invention and subsequent reconstitution. Preferably at least 40, 50, 60, 70, 80, 90, or 95% of the active agent's antigenicity is retained as assessed by ELISA.

Highly viscous liquids of the invention preferably comprise a glass forming polyol selected from the group consisting of glucose, maltulose, iso-maltulose, lactulose, sucrose, maltose, lactose, sorbitol, iso-maltose, maltitol, lactitol, palatinit, trehalose, raffinose, stachyose, melezitose and dextran.

5

Highly viscous liquid of the invention may contain any of the active agents described above. The active agent preserved by the highly viscous liquid may comprise a biological system, for instance cells, subcellular compositions, bacteria, outer membrane vesicle preparations and viruses. It may alternatively or further comprise a
10 molecules, for example proteins, peptides, amino acids, polynucleic acids, oligonucleotides, polysaccharides, oligosaccharides, polysaccharide – protein conjugates, oligosaccharide-protein conjugates. It may also comprise combinations of comprising two or more of the above active agents.

15 Preferred embodiments include a highly viscous liquid preferably obtained or obtainable by a method of the invention wherein the active agent is or comprises a vaccine or vaccine component. Preferred components of the vaccine are described above and include IPV, more preferably IPV and bacterial polysaccharides, preferably polysaccharides or oligosaccharides from *Haemophilus influenzae* b and *Neisseria*
20 *meningitidis* A, C, W and Y.

Preferred vaccine components include IPV (an inactivated mixture of polio virus strains). Preferably, IPV is combined with one or more of Hib PRP polysaccharide and/or meningococcal A, C, W and/or Y polysaccharides and/or pneumococcal
25 polysaccharides (as described above), more preferably IPV and Hib; IPV and MenC; IPV, Hib and MenC; Hib and MenC; IPV and MenA and C; Hib and Men A and C; IPV, Hib, Men A and C; Hib, Men C and Y; or IPV, Hib, Men C and Y.

In the above compositions where polysaccharides are used, oligosaccharides may also
30 be employed (as defined above).

Although these compositions may be adjuvanted (as described above), they are preferably unadjuvanted or preferably do not comprise aluminium salts.

5 Preferably the polysaccharides or oligosaccharides are conjugated to a peptide or carrier protein comprising T-helper epitopes (as described above).

The highly viscous liquid of the invention are preferably combined with other antigens in a combination vaccine which are optionally desiccated or preferably liquid formulations which can be used to reconstitute the dried components. Preferred
10 antigens to be combined with the contents of the container of the invention include one or more of diphtheria toxoid, tetanus toxoid, whole cell pertussis (Pw), acellular pertussis (Pa) (as described above), Hepatitis B surface antigen, pneumococcal polysaccharides, pneumococcal proteins, neisserial polysaccharides, neisserial proteins. Bacterial polysaccharides may be conjugated to a carrier protein such as
15 tetanus toxoid, tetanus toxoid fragment C, diphtheria toxoid, CRM197, pneumolysin, Protein D (US6342224) as described above.

A further aspect of the invention is a method of making a vaccine comprising the step of reconstituting the highly viscous liquid in an aqueous solution. In a preferred
20 embodiment, the aqueous solution comprises D, T and P (acellular or whole cell) vaccine. The DTP vaccine is optionally at least in part adjuvanted with aluminium hydroxide.

Another embodiment of the invention is a kit comprising the highly viscous liquid of
25 the invention held in a first container and a vaccine comprising liquid DTP (acellular or whole cell) in a second container.

All references or patent applications cited within this patent specification are incorporated by reference herein.

Examples

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1. Establishment of freezing conditions

Samples were made by dissolving sucrose in water to give 1%, 5%, 10% and 20% solutions. Samples were put into a Heto Drywinner 8-85 freeze dryer in which shelf temperature may be regulated to within 1°C, the final temperature of the condenser is -85°C, pressure is regulated with a bleed valve and 6 thermocouples are available to measure the product temperature. The shelf temperature setting was maintained at 15 °C throughout the process. The pressure was initially reduced to 200mBar and maintained at this level for 10 minutes before reducing the pressure further to 50mBars, 5mBars, 2.5mBars, 0.75mBars, 0.4mBars and 0.2mBars. Each pressure level was maintained for 20 minutes to allow the temperature to equilibrate and the temperature of the sample was read using a thermocouple. Thermocouples were attached to samples with different sucrose concentrations and the temperatures recorded in table 1 are mean values of the temperatures.

Results

All samples froze between 1.66 and 1.11mbars, irrespective of the concentration of sucrose present. The temperatures measured at different pressures were very close to those predicted from the triple point curve. Therefore the presence of sucrose does not have a large effect on the temperature of the samples at different pressures.

In order to avoid freezing of the sample, the pressure should be maintained above 2mBars for a shelf temperature of 15°C. At lower temperatures the pressure should be maintained at a higher level whereas use of a higher temperature would allow the pressure to be reduced further without the samples freezing.

Table 1

Pressure	Measured temperature	Theoretical temperature	Liquid/frozen
1000mBar	15 °C		liquid
50mBar	15 °C		liquid
5mBar	1 °C	1 °C	liquid
2.5mBar	-5 °C	-7 °C	liquid
0.75mBar	-21 °C	-21 °C	frozen
0.4mBar	-22 °C	-27 °C	frozen
0.2mBar	-27 °C	-32 °C	frozen

5 **Example 2. Method for drying without freezing or foam formation**

- Pr servation samples containing 5%, 10%, 15% and 25% sucrose were made and added to vials. Samples were put into a freeze dryer at a temperature setting of 15 °C throughout the process. The pressure was initially reduced to 200mBars and
- 10 maintained at this level for 10 minutes to allow degassing before reducing the pressure further. The pressure was further reduced to 8mbars for two to three hours during which time thermocouples inside the samples showed that the sample temperature reduced to 4 °C due to evaporative cooling. After 2-3 hours, the temperature of the samples returned to 15 °C; indicating that evaporation under these
- 15 temperature and pressure conditions was near completion. During this stage of the process, the sample did not boil to form a foam or freeze so that an active agent within the sample is exposed to as little stress as possible. The sample have the appearance of viscous liquid.
- 20 Further drying of the samples was achieved by reducing the pressure further to 0.1mbars while keeping the shelf temperature setting at 15 °C. These conditions were maintained for a further 10-16 hours. During this phase, the sample temperature remained at 15 °C since the rate of evaporation was slow. Further drying took place

and the resultant sample had a solid appearance. If the sample was placed on its side, the sample contents flowed very slowly, over a period of days showing that the sample is a liquid glass of high viscosity. Figure 1 shows the appearance of the high viscosity liquid.

5

Example 3 Retention of IPV immunogenicity after drying without freezing or foam formation

Such samples have not been subjected to stresses associated with the bubbling that accompanies foam formation or freezing. Experiments were performed to determine whether this method produced a high level of antigen retention when used to dry IPV.

Three separate experiments were performed in which IPV was resuspended in an aqueous solution with 10% sucrose or 10% trehalose as the stabilising agent. The samples were put into siliconised vials which were placed into a Heto Drywinner 8-85 freeze-dryer and the temperature was set to 15 °C. The pressure was initially reduced to 35mBars to degas the sample. After 10 minutes, the pressure was further reduced to 8mBars and was kept at this level for two hours. During this period the temperature setting was kept at 15 °C and the temperature into the sample was monitored. As water evaporated from the sample, the temperature dropped to 4 °C but towards the end of the two hours, the temperature returned to 15 °C as the rate of evaporation slowed. No bubbling or foam formation occurred under these conditions. The pressure was then reduced further to 0.1mbars and these conditions were maintained for 16 hours more in the first two experiments and for 10 hours more in the third experiment.

25

The samples were reconstituted in water and an ELISA was used to assess the integrity of structure of the three polio virus strains. The monoclonal antibody against type 3 IPV, was used in an ELISA to assess the degree of antigen retention in the reconstituted, freeze dried sample compared to a reference sample that had not been frozen. Results are presented as a percentage of the reading given for a sample which had not undergone a drying procedure.

30

Results

The dried samples had a solid appearance however they appeared to be in the form of a highly viscous liquid/glass since, over a period of days, the dried sample was able to flow if the container was inverted.

5

Table 2 Retention of type 3 IPV antigen as determined by ELISA using a monoclonal antibody (drying without foaming or freezing)

Formulation	1 st experiment (18 hour cycle)	2 nd experiment (18 hour cycle)	3 rd experiment (12 hour cycle)
10% sucrose	75%	78%	91%
10% trehalose	82%	79%	93%

10 These levels of type 3 IPV antigen retention compares very favourably with the foam drying and freeze drying results shown below where very low values were usually found in the same ELISA format when a monoclonal antibody against type 3 was used .

15 Table 3 Retention of type 1, 2 and 3 IPV antigens as determined by ELISA using a monoclonal and polyclonal antibodies (freeze drying and foam drying)

Method of drying	Polyol content	ELISA – type 1/2/3 %	
		Polyclonal	Monoclonal
Freeze drying	3.15% sucrose	46/49/58*	25/0/0
Foam drying	3.15% sucrose	85/97/106	55/68/57
Freeze drying	10% trehalose	47/43/58	
Foam drying	10% trehalose	93/86/84	72/75/87

* The experiment freeze drying in the presence of 3.15% sucrose was repeated five
20 times and the results shown are from one representative experiment.

Example 4 Long term storage stability of dried IPV stored as a highly viscous liquid/glass.

IPV dried using the method described in Example 3 was stored at 4 °C for 9 months.

- 5 The samples were reconstituted in water and an ELISA was used to assess the integrity of structure of the three polio virus strains. Three monoclonal antibodies, one against each strain, were used in separate ELISAs to assess the degree of antigen retention in the reconstituted stored sample. A similar ELISA had been carried out on reconstituted samples from the same batch prior to storage. All results were compared
- 10 to a reference sample that had not been dried. Results are presented as a percentage of the reading given for a sample which had not undergone a drying procedure.

Results

- 15 Table 4. Retention of IPV antigens after storage as a highly viscous liquid for 9 months

Treatment	Type 1 ELISA	Type 2 ELISA	Type 3 ELISA
Dried/reconstituted Not stored	72%	75%	88%
Dried/reconstituted 9 months 4 °C	70%	94%	90%

- 20 Therefore IPV which has been dried by the method described in Example 3 can be stored at 4 °C for at least 9 months without loss of antigenicity.

Claims

1. A method for preserving an active agent comprising the steps of:
 - a) preparing a preservation sample by dissolving/suspending an active agent in a solution of a stabilising agent;
 - b) subjecting the preservation sample to such temperature and pressure conditions so that the preservation sample loses solvent by evaporation, without freezing or bubbling involved in foam formation, to form a viscous liquid;
 - c) removing solvent until the preservation sample dries to form a highly viscous liquid.
2. The method of claim 1 wherein the pressure is reduced to 20 mbars or below during step b).
3. The method of claim 1-2 wherein the temperature external to the preservation sample is between 5°C and 37°C during step b).
4. The method of claim 3 wherein the temperature external to the preservation sample is between 5°C and 37°C during step c).
5. The method of claim 1-4 wherein the temperature external to the preservation sample is higher during step c) than it is in step b).
6. The method of claim 5 wherein the temperature external to the preservation sample is increased to above 20°C during step c).
7. The method of claim 1-6 wherein the pressure is reduced in step c) compared to the pressure during step b).
8. The method of claim 7 wherein the pressure is reduced to 1mbar or below during step c).

9. The method of claim 1-8 wherein steps b) and c) are completed in less than 12 hours.
10. The method of claims 1-9 wherein the stabilising agent comprises a glass forming polyol, selected from the group consisting of glucose, maltulose, iso-maltulose, lactulose, sucrose, maltose, lactose, sorbitol, iso-maltose, maltitol, lactitol, palatinit, trehalose, raffinose, stachyose, melezitose and dextran.
11. The method of claim 10 wherein the concentration of stabilising agent is less than 15%.
12. The method of claims 1-11 wherein the preservation sample is dried in a container with a solvent repellent interior surface.
13. The method of claims 1-12 wherein the active agent comprises a molecule selected from the group consisting of protein, peptide, amino acid, polynucleotide, oligonucleotide, polysaccharide, oligosaccharide, polysaccharide-protein conjugate and oligosaccharide-protein conjugate.
14. The method of claim 1-12 wherein the active agent comprises a biological system selected from the group consisting of cells, subcellular compositions, bacteria, viruses, virus components and virus like particles.
15. The method of claim 14 wherein the active agent comprises IPV (inactivated polio virus).
16. The method of claim 14-15 wherein the active agent comprises Hib (*Haemophilus influenzae* type b) polysaccharide or oligosaccharide.
17. The method of claim 14-16 wherein the active agent comprises *Neisseria meningitidis* C polysaccharide or oligosaccharide.

18. The method of claims 1-17 wherein the active agent comprises a vaccine.
19. A highly viscous liquid comprising an active agent wherein the antigenicity or activity of the active agent is preserved.
- 5 20. The highly viscous liquid of claim 19 obtainable by the method of claims 1-18.
21. The highly viscous liquid of claim 19 or 20 comprising a glass forming polyol selected from the group consisting of glucose, maltulose, iso-maltulose, lactulose, sucrose, maltose, lactose, sorbitol, iso-maltose, maltitol, lactitol, palatinit, 10 trehalose, raffinose, stachyose, melezitose and dextran.
22. The highly viscous liquid of claim 19-21 wherein the active agent comprises comprises a molecule selected from the group consisting of protein, peptide, amino acid, polynucleotide, oligonucleotide, polysaccharide, oligosaccharide, polysaccharide-protein conjugate and oligosaccharide-protein conjugate.
- 15 23. The highly viscous liquid of claim 19-21 wherein the active agent comprises a biological system selected from the group consisting of cells, subcellular compositions, bacteria, viruses, virus components and virus like particles.
- 20 24. The highly viscous liquid of claim 19-23 wherein the active agent comprises a vaccine.
- 25 25. The highly viscous liquid of claim 19-24 wherein the active agent comprises IPV.
26. The highly viscous liquid of claim 19-25 wherein the active agent comprises IPV and a bacterial polysaccharide.
- 30 27. The highly viscous liquid of claim 19-26 wherein the active agent comprises Hib (*Haemophilus influenzae* b) polysaccharide or oligosaccharide, preferably conjugated to a carrier protein.

28. The highly viscous liquid of claim 19-27 wherein the active agent comprises
Neisseria meningitidis serogroup C polysaccharide or oligosaccharide, preferably
conjugated to a carrier protein.

5

29. The highly viscous liquid of claim 19-28 held within a container with a solvent
repellent interior surface.

10

30. An immunogenic composition or vaccine comprising the highly viscous liquid of
claim 19-29 and a pharmaceutically acceptable excipient.

31. A method of making a vaccine comprising the step of reconstituting the highly
viscous liquid of claim 19-29 in an aqueous solution.

15

32. The method of claim 31 wherein the aqueous solution comprises D, T and P
(acellular or whole cell) vaccine.

33. The method of claim 32 where the DTP vaccine is at least in part adjuvanted with
aluminium hydroxide.

20

34. A kit comprising the highly viscous liquid of claims 19-29 held in a first container
and liquid DTP (acellular or whole cell) vaccine in a second container.

1/1

Figure 1

